

Recent Results From EBIT-II Using A Spare Astro-E Microcalorimeter

G. V. Brown, E. Behar, P. Beiersdorfer, K. R. Boyce, H. Chen, K. C. Gendreau, J. Gygax, S. M. Kahn, R. L. Kelley, F. S. Porter, C. K. Stahle, A. E. Szymkowiak

This article was submitted to
New Century of X-Ray Astronomy
Yokohama, Japan
March 6-8, 2001

U.S. Department of Energy

Lawrence
Livermore
National
Laboratory

June 26, 2001

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This report has been reproduced
directly from the best available copy.

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information
P.O. Box 62, Oak Ridge, TN 37831
Prices available from (423) 576-8401
<http://apollo.osti.gov/bridge/>

Available to the public from the
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Rd.,
Springfield, VA 22161
<http://www.ntis.gov/>

OR

Lawrence Livermore National Laboratory
Technical Information Department's Digital Library
<http://www.llnl.gov/tid/Library.html>

Recent results from EBIT-II using a spare Astro-E microcalorimeter

G. V. Brown¹, E. Behar³, P. Beiersdorfer², K. R. Boyce¹, H. Chen², K. C. Gendreau¹, J. Gygax¹, S. M. Kahn³, R. L. Kelley¹, F. S. Porter¹, C. K. Stahle¹, and A. E. Szymkowiak¹

¹NASA/GSFC, Greenbelt, Maryland, USA

²Lawrence Livermore National Laboratory, Livermore, CA, USA

³Columbia Astrophysics Laboratory, Columbia University, New York, NY, USA

ABSTRACT

A spare NASA/GSFC Astro-E microcalorimeter has been installed, tested, and run successfully on EBIT-II at the Lawrence Livermore National Laboratory. A brief overview of results including measurements by the microcalorimeter of absolute excitation cross sections, time dependent spectra, and spectra as a function of Maxwellian temperature are discussed.

Over the last decade the spectroscopy group at the Lawrence Livermore National Laboratory electron beam ion trap EBIT-II, in collaboration with the Columbia University Astrophysics Laboratory, and recently, NASA's Goddard Space Flight Center, has developed an extensive laboratory astrophysics program. Recently, a spare NASA/GSFC *Astro-E* XRS microcalorimeter has been installed, tested, and run successfully on EBIT-II (Porter et al. 2000). The microcalorimeter complements crystal and grating spectrometers making it possible to measure a broad band-width ($\sim 0.3 - 10$ keV) with moderate resolution while simultaneously measuring a narrow bandwidth ($\sim 0.7 - 1.3$ keV) with high resolution. In conjunction with the spectrometers already present at EBIT-II, the microcalorimeter has been used to measure absolute excitation cross sections, time dependent spectra, and spectra as a function of Maxwellian temperature.

The broad band-width and ~ 10 eV resolution of the XRS enables us to normalize cross sections to the radiative recombination (RR) photon emission. Because radiative recombination is the simplest atomic scattering process, highly-accurate RR cross sections are known for high electron-impact energies. Once normalized to the RR, our measured cross sections are, therefore, “absolute” in the sense that they are normalized to the most-highly accurate and independent process available. Some of the first results from these measurement can be found in Chen et al., 2001 and in Porter et al. 2001.

Using the ability to measure time dependent spectra with the XRS, we have studied ionizing plasmas, i.e., plasmas whose charge balance has not yet reached equilibrium. Figure 1 shows two spectra measured at different times: 1(a) is near equilibrium, and 1(b) is before equilibrium is reached. This shows that the ratio of $3s \rightarrow 2p$ to $3d \rightarrow 2p$ lines in Fe XVII changes dramatically between the ionizing and the equilibrium condition. It is before equilibrium is reached, i.e., during ionization, that processes such as innershell ionization and excitation play an important role in the spectra. Our ability to measure time resolved spectra makes it possible to accurately discriminate between lines emitted during ionization from those emitted during equilibrium.

We have presented a brief overview of work being done as part of the Laboratory Astrophysics program at the LLNL EBIT-II. Other work includes using the Maxwellian simulation mode of EBIT-II to create and measure spectra emitted from plasmas at specific electron temperatures to determine the correct charge balance for a given Maxwellian temperature, and measuring the relative intensity of O VIII Lyman α to Lyman β (Gendreau et al. 2001). Laboratory data of these types will make the analysis of astrophysical spectra more reliable.

This work was performed while G. V. Brown held a NRC Research Associateship award at GSFC, and under the auspices of the U.S. D.o.E. by U. of California LLNL under contract No. W-7405-Eng-48 and supported by NASA SARA grants to LLNL, GSFC, and Columbia University.

REFERENCES

- Chen, H., et al. J. 2001, *Astrophys. J.*, in prep
- Gendreau, K. C., et al. 2001, *Astrophys. J.*, in prep
- Porter, F. S., et al. 2000, in *Proceedings of the 45th annual SPIE meeting on Optical Science and Technology* (SPIE Press), in press
- Porter, F. S., et al. 2001, in *Proceedings of a New Century of X-ray Astronomy*




Fig. 1.— (a) Iron spectrum measured with the XRS microcalorimeter measured near equilibrium for a quasi-Maxwellian energy distribution. (b) Same spectrum as (a) during ionization. Notice the large difference between the $3s \rightarrow 2p$ and $3d \rightarrow 2p$ lines in Fe XVII.

University of California
Lawrence Livermore National Laboratory
Technical Information Department
Livermore, CA 94551